

Nursing Home Quality Measures(Q3) VA-AVG

Based on us-avg-q3

State	Percent of long-stay residents given influenza vaccination during the flu season
VA	92.2840909090909%

Nursing Home Quality Measures(Q3) VA-AVG

Based on us-avg-q3

Percent of long-stay residents who were assessed and given pneumococcal
vaccination

90.49438202247191%

Nursing Home Quality Measures(Q3) VA-AVG

Based on us-avg-q3

Percent of long-stay residents whose need for help with daily activities has increased



16.607547169811321%

Nursing Home Quality Measures(Q3) VA-AVG

Based on us-avg-q3

Percent of high-risk long-stay residents who have pressure sores



11.259398496240602%

Nursing Home Quality Measures(Q3) VA-AVG

Based on us-avg-q3

Percent of low-risk long-stay residents who have pressure sores



2.284403669724771%

Nursing Home Quality Measures(Q3) VA-AVG

Based on us-avg-q3

Percent of low-risk long-stay residents who lose control of their bowels or bladder

58.509505703422053%

Nursing Home Quality Measures(Q3) VA-AVG

Based on us-avg-q3

Percent of long-stay residents who have/had a catheter inserted and left in their bladder



4.112781954887218%

Nursing Home Quality Measures(Q3) VA-AVG

Based on us-avg-q3

Percent of long-stay residents who had a urinary tract infection



10.254681647940075%

Nursing Home Quality Measures(Q3) VA-AVG

Based on us-avg-q3

Percent of long-stay residents who lose too much weight



8.970037453183521%

Nursing Home Quality Measures(Q3) VA-AVG

Based on us-avg-q3

Percent of short-stay residents given influenza vaccination during the flu season

82.490494296577947%

Nursing Home Quality Measures(Q3) VA-AVG

Based on us-avg-q3

Percent of short-stay residents who have pressure sores



12.140151515151515%